

CPD Answer form

Please **PRINT** your details below:

Title

First Name

Last Name

Address

.....

.....

City

County

Postal Code

E-mail

Your Work

What is the name of your practice/
CDS/hospital?

.....

What is your job title?

.....

Please **TICK** (✓) the answer to
each question

Q.1 A B C D

Q.2 A B C D

Q.3 A B C D


Q.4 A B C D

If you prefer not to be contacted for purposes other
than this competition, please tick this box. We will not
pass your details on to any third parties.

Please send your completed
answer form to:
CPD Questions, *Vital* magazine,
BDJ, Nature Publishing Group,
4-6 Crinan Street,
London N1 9XW.

CPD Questions

From summer 2008, *Vital* will be offering verifiable CPD when it becomes mandatory for DCPs. In the meantime we have provided some CPD style questions for our 'Vital Guide to...' series to encourage you to anticipate future CPD requirements. By way of extra encouragement we are pleased to offer a copy of *Management of medical emergencies for the dental team* to the senders of the first three correct entries to be drawn on 2 May 2008. The answers to the questions will be published in the next issue of *Vital*. After reading the article have a go at the following questions and send in your completed form to us. In the context of this article, only one of the answers to each of the following questions is correct.



- In England and Wales how many patients under 18-years-old have fixed braces fitted under the NHS every year?
 - Under 130,000
 - Over 130,000
 - Approximately 13,000
 - Approximately 18,000
- According to the IOTN, approximately what percentage of 11 to 12-year-old children in the UK would benefit from orthodontic treatment?
 - 19-25%
 - 33%
 - 35%
 - 11-12%
- Which of the following might be carried out as an element of orthodontic treatment?
 - Removal of a deciduous canine to allow spontaneous correction of the centreline
 - Removal of a deciduous canine where the permanent unerupted canine is becoming ectopically positioned
 - The use of functional appliances to reduce an increased overjet and encourage the lower jaw to come forwards
 - All of the above
- Which of the following is incorrect?
 - Between 0.4% and 1% of incisor teeth lose a third or more of their root length during orthodontic treatment
 - Orthodontic patients are at risk of gingivitis and periodontal disease
 - Orthodontic therapists are not yet recognised within the UK
 - Dental students are taught to perform a straightforward orthodontic diagnosis